



MEASURES TO PREVENT
RUGOSE TOMATO VIRUS

PRODUCTION AREA RECOMMENDATIONS




1

PRODUCTION AREA RECOMMENDATIONS:



Cutting tools can be a factor in disease spread. Disinfect them periodically during the workday, whenever changing greenhouses, and at the end of the day.



2

PRODUCTION AREA RECOMMENDATIONS:



Allow entry into the production area only for essential personnel.

3

PRODUCTION AREA RECOMMENDATIONS:

Ensure that anyone entering the production area complies with self-hygiene recommendations. Provide enough coveralls, gowns, and disposable gloves for the entire team.

4

PRODUCTION AREA RECOMMENDATIONS:

Place effective disinfectant footbaths at all entrances, for use by people and machines.



5

PRODUCTION AREA RECOMMENDATIONS:



Follow good hygiene practices: clean work cabins with disinfectant products and sanitize tools.

6

PRODUCTION AREA RECOMMENDATIONS:



Avoid taking objects from the inside of the cultivation area outdoors.



7

PRODUCTION AREA RECOMMENDATIONS:



Never park vehicles near the cultivation area to prevent dispersion with tires.

For more information, visit:

[**www.semillasfito.com/protectagainstrugose**](http://www.semillasfito.com/protectagainstrugose)

improving
food is in our
genes



semillasfito.com

